

6042 Cornerstone Court West, Suite B

San Diego, CA 92121 Tel: 1.858.829.3082 Fax: 1.858.481.8694

Email: info@bpsbioscience.com

Data Sheet

JARID1C (KDM5C)

Human recombinant, with N-terminal FLAG-tag and His-tag

Catalog #: 50112

Lot #: 140709 **Conc.** 1.3 mg/ml

Formulated in: 40 mM Tris-HCl, 110 mM NaCl, 2.2 mM KCl, 0.04% Tween-20, 80 μg/mL and 20% glycerol.

Stability: At least 6 months at -80°C. Avoid freeze/thaw cycles. Storing diluted enzyme is not recommended, if necessary, use carrier protein (BSA 0.1 - 0.5%).

References:

- 1. Xu, J., et al. (2008). PLoS One **3(7)**:e2553.
- 2. Iwase, S., et al. (2007). Cell **128(6):**1077-88.


<u>Description</u>: Human JARID1C, also known as SMCX (Smcy homolog, X-linked), KDM5C, or XE169, GenBank Accession No. NM_004187), full length [a.a. 2-1560(end)] with N-terminal FLAG- and His-tags, MW= 181 kDa, expressed in Sf9 cells via a baculovirus expression system.


Specific Activity: 0.14 pmol/min/µg.


Assay Conditions: Assay was conducted using AlphaScreen® technology using BPS JARID1C Homogeneous Assay Kit, Catalog #50511. JARID1C enzyme is incubated with biotinylated histone H3 peptide substrate in JARID assay buffer in a 10 ul reaction for 2 hours at RT. AlphaLISA® anti-Mouse Acceptor beads (Perkin Elmer) are added,followed by the primary antibody addition and 30 min incubation at RT. AlphaScreen® Streptavidin-conjugated donor beads (Perkin Elmer) are added. After 10-15 minute incubation at room temperature, Alphacounts are measured.

Application: Useful for the study of enzyme kinetics, screening inhibitors, and selectivity profiling.

Quality Assurance







OUR PRODUCTS ARE FOR RESEARCH USE ONLY. NOT FOR DIAGNOSTIC OR THERAPEUTIC USE.